

Oct 28th, 2:30 PM - 4:00 PM

# Cultivation of the Potato in Kosovo and Analysis of Some Quality Parameters

Fidan Feka

*University for Business and Technology*

Shkëlzim Ukaj

*University for Business and Technology, shkelzim.ukaj@ubt-uni.net*

Syzana Aliu

*University for Business and Technology*

Fjolla Vllasaliu

*University for Business and Technology*

Sara Selimi

*University for Business and Technology*

*See next page for additional authors*

Follow this and additional works at: <https://knowledgecenter.ubt-uni.net/conference>

 Part of the [Food Science Commons](#)

---

## Recommended Citation

Feka, Fidan; Ukaj, Shkëlzim; Aliu, Syzana; Vllasaliu, Fjolla; Selimi, Sara; Vllasaliu, Taulant; and Bruqi, Era, "Cultivation of the Potato in Kosovo and Analysis of Some Quality Parameters" (2017). *UBT International Conference*. 152.

<https://knowledgecenter.ubt-uni.net/conference/2017/all-events/152>

This Event is brought to you for free and open access by the Publication and Journals at UBT Knowledge Center. It has been accepted for inclusion in UBT International Conference by an authorized administrator of UBT Knowledge Center. For more information, please contact [knowledge.center@ubt-uni.net](mailto:knowledge.center@ubt-uni.net).

---

**Presenter Information**

Fidan Feka, Shkëlzim Ukaj, Syzana Aliu, Fjolla Vllasaliu, Sara Selimi, Taulant Vllasaliu, and Era Bruqi

# CULTIVATION OF THE POTATO IN KOSOVO AND ANALYSIS OF SOME QUALITY PARAMETERS

Fidan Feka, Shkëlzim Ukaj, Syzana Aliu, Fjolla Vllasaliu, Sara Selimi, Taulant Vllasaliu, Era Bruqi

UBT – Higher Education Institution, Lagjja Kalabria, 10000 p.n., Prishtine,  
Kosovo  
shkelzim.ukaj@ubt-uni.net

**Abstract.** Potatoes in Kosovo are cultivated in about 10,000 ha with a low average production of 10-12 t /ha and represent one of the main agricultural crops, both for agro-economic, physiological and productive importance, as well as in feeding the population. The purpose of the study was to research the level of potato cultivation in Kosovo and to determine some toxic elements of potato cultivated in some localities of Kosovo and comparing this level with the standards allowed by the World Health Organization regulation ,also comparing it with the standards allowed by Codex Stan 193-1995, for fruits and vegetables. Based on the results obtained for elements Zn, Cu, Pb and Cd , there was an increase of heavy metal concentrations in potato compared to the level allowed by the World Health Organization (WHO).

**Keywords:** Potato cultivation in Kosovo, potato quality and its importance.

## INTRODUCTION

Patato is a wide spread plant, with high production skills and multifunctional usage. It is a daily component in feeding people, the first subject and very important in the food processing industry, in which is created starch, alcohol and other products.<sup>[1]</sup> The chemical composition of the patato is complex enough and it depends from the cultivator, the conditions of the cultivation, manner of feeding, cultivation areas etc. In average the chemical composition of the patato goes around 75% of water, 17-23% starch, 2% nitrogen compounds and 1% mineral materials, 1% cellulose, 0.15% grease, 0.4% sugar, 2.5% pectin, significant amounts of C and B vitamins, various acids of (apple, lemon, wine) and various enzymes. <sup>[2],[3]</sup>



**Fig. 1.** Patato

Earth is one of the most important resources of the contamination of plant foods from heavy metals that enter in the human and animal organism through the food chain.

The term “heavy metals” refers to each metal element that has a relatively high density, which is toxic or poisonous, and in high concentration. Heavy metals are created in the top layer of the earth and can be transferred from the earth in other components of ecosystem, like in underground waters or in harvest, and can also influence in the health of people through the supply of water and food chain. High concentrations of Pb and Cd in diets are linked with the development of many diseases, especially those cardiovascular, kidney, nervous and ossuary tissues. [4],[5]

The cadmium accumulation (Cd) in the human body can cause injuries of kidney, bones and lungs, whereas the accumulation of bullet may damage the central nervous system, kidney and blood system.

## **MATERIAL AND METHODS**

The identification of the area for getting the samples was based in those locations where we thought we may have influence from outside factors, and we were mainly oriented in those spaces. The gained samples were taken in the region of Vushtrri and Obiliq.<sup>[6]</sup>

Primarily, the gained samples were stuffed in small particles and then they were grinded. From the grinded mass were taken 20g and were placed in the mineralization tank, treating it in 400°C temperature in 60 min.



The liquid mass was filtered and then the toxic elements were analyzed through Spectrophotometer HACH 2000.


## **RESULTS AND DISCUSSION**

From the analyzed results for Cd, Pb, Zn and Cu elements, we can conclude that in gained patatoe samples we have an overrun concentration of Zn, expressed in mg/kg, conforming to the prescribed standard from The World Organization of Health.

Also, in the locations Vushtri/ Akrashtice we have and overrun of the standard, seen in the first figure of distribution of copper.

In Vushtri we have had a fixed concentration of elements comparing to the allowed concentration from WHO. This increase has come as a results of pollution of the river Sitnica and from the resident who has had planted potatoes and watered them directly with the same water.





**Figure 2.** Heavy metal content (mg/kg) by localities.

## CONCLUSIONS

From the gained results we can conclude that:

- The irrigation of agricultural lands should be strictly prohibited in the river Sitnica, where this phenomenon is still applied in some localities. In the most of spaces this phenomenon is not applied, as a result of the functionalization of the water system.
- Also, a suggestion for the residents that cultivate this product and others should analyze the agricultural lands before having a realistic mirror.
- Should make systematic analysis of authorized people to this.
- Especially, the increase of concentration level of Pb, can cause highlighted consequences to children.

## REFERENCES

1. Cheraghi M., Lorestani B., Merrikhpour H. & Rouniasi N., Heavy metal risk assessment for Potatoes grown in over used phosphate fertilized soils, *Environ. Monit. Assess.* 185(2), 1825-1831, 2013.
2. CODEX STAN 193-195., Codex General Standard for Contaminants and Toxins in Food and Feed. Adopted 1995, Amended 2010. p 1, 26, 27.
3. Demirezen D. & Aksoy A., Heavy metal levels in vegetables in Turkey are within safe limits for Cu, Zn, Ni and exceeded for Cd and Pb, *Journal of Food Quality.* vol. 29, pp 252-265, 2006.
4. Ivana Maksimović., Marina Putnik-Delić., Ilin Ž. & Mirosavljević M., Essential (Cu and Zn) and trace (Pb and Cd) heavy metal loads in onion and potato. University of Banjaluka. *Agroznanje*, vol. 13, br.1. pp 113-122, 2012.
5. Janette Musilova., Judita Bystricka., Alena Vollmannova., Beata Volnova. & Alzbeta Hegedusova., Factors affecting heavy metals accumulation in potato tubers. *Environmental Protection and Natural Resources.* Vol. 26 No 3(65), pp 54-59, 2015.
6. Jorhem L. & Sundstroem B., Levels of Lead, Cadmium, Zinc, Copper, Nickel, Chromium, Manganese and Cobalt in foods on the Swedish market, *J. Food Comp. Anal.*, 6, pp 223-241, 1993.  
[http://www.mbpzhr-ks.net/repository/docs/TREGU\\_I\\_PATATES\\_FINAL.pdf](http://www.mbpzhr-ks.net/repository/docs/TREGU_I_PATATES_FINAL.pdf)