

University for Business and Technology in Kosovo

UBT Knowledge Center

UBT International Conference

2015 UBT International Conference

Nov 7th, 9:00 AM - 5:00 PM

New integral transform in Caputo type fractional difference operator

Artion Kashuri

University of Vlora, artionkashuri@gmail.com

Akli Fundo

Polytechnic University of Tirana, rozanaliko86@gmail.com

Rozana Liko

University of Vlora, aklifundo@yahoo.com

Follow this and additional works at: <https://knowledgecenter.ubt-uni.net/conference>



Part of the [Computer Sciences Commons](#), and the [Digital Communications and Networking Commons](#)

Recommended Citation

Kashuri, Artion; Fundo, Akli; and Liko, Rozana, "New integral transform in Caputo type fractional difference operator" (2015). *UBT International Conference*. 84.

<https://knowledgecenter.ubt-uni.net/conference/2015/all-events/84>

This Event is brought to you for free and open access by the Publication and Journals at UBT Knowledge Center. It has been accepted for inclusion in UBT International Conference by an authorized administrator of UBT Knowledge Center. For more information, please contact knowledge.center@ubt-uni.net.

New integral transform in Caputo type fractional difference operator

Artion Kashuri¹, Akli Fundo², Rozana Liko¹

¹Department of Mathematics, Faculty of Technic Sciences, University "Ismail Qemali", Vlora, Albania

²Department of Mathematics, Polytechnic University of Tirana, Albania
{artionkashuri¹, rozanaliko86²}@gmail.com, aklifundo@yahoo.com³

Abstract. In this paper, we introduce Caputo type nabla (q, h) -fractional difference operators and investigate their basic properties and also to show the applicability of this interesting (q, h) - new integral transform method and its efficiency in solving linear fractional difference equations. Differential equations with fractional derivative provide a natural framework for the discussion of various kinds of real problems modeled by the aid of fractional derivative.

Discrete analogues of some topics of continuous fractional calculus have been developed. Finally, we provide the general solutions in terms of discrete Mittag-Leffler functions.

Keywords: (q, h) - new integral transform, convolution, fractional difference equations, nabla (q, h) -fractional integral, nabla (q, h) - fractional derivative.

1. Introduction

(q, h) –new integral transform is derived from the classical Fourier integral and was introduced by PhD student Artion Kashuri and Associate Professor Akli Fundo to facilitate the process of solving linear fractional difference equations in the time domain [6], [7]. (q, h) –new integral transform is defined for functions of exponential order. Fractional calculus deals with the study of fractional order integrals and derivatives and their applications [1]. Riemann–Liouville and Caputo are kinds of fractional derivatives which generalize the ordinary integral and differential operators. Differential equations with fractional derivative provided a natural framework for the discussion of various kinds of real problems modeled by the aid of fractional derivative. Discrete analogues of some topics of continuous fractional calculus have been developed. The aim of this paper is to introduce Caputo type nabla (q, h) –fractional difference operators and investigate their basic properties, and also to show the applicability of this interesting (q, h) –new integral transform method and its efficiency in solving linear fractional difference equations. Here, we solve some linear fractional difference equations involving Caputo type (q, h) –derivatives and provide the general solutions in terms of discrete Mittag–Leffler functions.

2. Preliminaries

For the convenience of readers, we provide some basic concepts concerning the nabla calculus on time scales. By a time scale \mathbb{T} we understand any nonempty, closed subset of reals with the ordering inherited from reals. Thus the reals \mathbb{R} , the integers \mathbb{Z} , the natural numbers \mathbb{N} , the nonnegative integers \mathbb{N}_0 , the h –numbers $h\mathbb{Z} = \{hk: k \in \mathbb{Z}\}$ with fixed $h > 0$, and the q –numbers $q^{\mathbb{N}_0} = \{q^k: k \in \mathbb{N}_0\}$ with fixed $q > 1$ are examples of time scales. For any $t \in \mathbb{T}$, we define the forward and backward jump operators as $\sigma(t) := \inf\{s \in \mathbb{T}: s > t\}$ and $\rho(t) := \sup\{s \in \mathbb{T}: s < t\}$, respectively. The

forward and backward graininess functions are defined as $\mu(t) := \sigma(t) - t$ and $\mu(t) := t - \rho(t)$, respectively. By convention, $\inf\{\emptyset\} = \sup\{\mathbb{T}\}$ and $\sup\{\emptyset\} = \inf\{\mathbb{T}\}$ for these operators. We say that a point is left-dense if $\rho(t) = t$ and left-scattered if $\rho(t) \neq t$. The right-dense and right-scattered points are defined in a similar manner. A point which is both left- and right-scattered is discrete. If $\inf\{\mathbb{T}\} = a_0 > -\infty$, we define $\mathbb{T}_k := \mathbb{T} \setminus a_0$, otherwise $\mathbb{T}_k := \mathbb{T}$. For a function $f: \mathbb{T} \rightarrow \mathbb{R}$ and a point $t \in \mathbb{T}_k$, we define $f^\nabla(t)$ to be a number such that for $\epsilon > 0$ there exists a neighborhood $\mathcal{U} \subset \mathbb{T}$ of t which satisfies

$$|f(\rho(t)) - f(\tau) - f^\nabla(t)[\tau - \rho(t)]| \leq \epsilon|\tau - \rho(t)| \text{ for all } \tau \in \mathcal{U}.$$

If $f^\nabla(t)$ is defined for all $t \in \mathbb{T}_k$, then the function obtained is called the ∇ -derivative of f . A function $f: \mathbb{T} \rightarrow \mathbb{R}$ is called left-dense continuous or ld-continuous provided it is continuous at every left-dense point in \mathbb{T} , and $f(t+)$ exists for every right-dense point in \mathbb{T} , and the set of ld-continuous functions is denoted by $C_{ld}(\mathbb{T})$. If $f \in C_{ld}(\mathbb{T})$, then there is a function F such that $F^\nabla(t) = f(t)$. In this case, we define the ∇ -integral as

$$\int_a^t f(\tau) \nabla\tau = F(t) - F(a) \text{ for all } t \in \mathbb{T}.$$

Clearly, if $f: \mathbb{T} \rightarrow \mathbb{R}$ is ld-continuous and $t \in \mathbb{T}_k$, then

$$\int_{\rho(t)}^t f(\tau) \nabla\tau = v(t)f(t).$$

A function $f \in C_{ld}(\mathbb{T})$ is called v -regressive if $1 + fv \neq 0$ on \mathbb{T}_k , and $f \in C_{ld}(\mathbb{T})$ is called positively- v -regressive if $1 + fv > 0$ on \mathbb{T}_k . The set of v -regressive functions and the set of positively v -regressive functions are denoted by $\mathcal{R}_v(\mathbb{T})$ and $\mathcal{R}_v^+(\mathbb{T})$, respectively. For simplicity, we denote by ${}_c\mathcal{R}_v(\mathbb{T})$ the set of v -regressive constants. Let $\lambda \in {}_c\mathcal{R}_v(\mathbb{T})$ and $s \in \mathbb{T}$, then the generalized exponential function $\hat{e}_\lambda(\cdot, s)$ on time scale \mathbb{T} is denoted by the unique solution of the initial value problem

$$\begin{cases} x^\nabla(t) = \lambda x(t), & t \in \mathbb{T}_k \\ x(s) = 1. \end{cases}$$

For $p \in {}_c\mathcal{R}_v(\mathbb{T})$, define circle minus p by

$$\ominus_v p := -\frac{p}{1 - pv}.$$

Then the unique solution of the initial value problem

$$\begin{cases} x^\nabla(t) = -\lambda x(t), & t \in \mathbb{T}_k \\ x(s) = 1. \end{cases}$$

takes the form $\hat{e}_{\ominus_v \lambda}(\cdot, s)$. It is known that the exponential function $\hat{e}_f(\cdot, s)$ is strictly positive on $[s, \infty[_\mathbb{T}$ provided that $f \in \mathcal{R}^+([s, \infty[_\mathbb{T})$. The definition of the generalized monomials $\hat{h}_n: \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}$ ($n \in \mathbb{N}_0$) is given by

$$\hat{h}_n(t, s) = \begin{cases} 1, & n = 0 \\ \int_s^t \hat{h}_{n-1}(\tau, s) \nabla\tau, & n \in \mathbb{N}. \end{cases}$$

for $s, t \in \mathbb{T}$, [2]. If we let $\hat{h}_n^\nabla(t, s)$ denote for each fixed $s \in \mathbb{T}$ the derivatives of $\hat{h}_n^\nabla(t, s)$ with respect to t , then

$$\hat{h}_n^\nabla(t, s) = \hat{h}_{n-1}(t, s) \text{ for } n \in \mathbb{N}, t \in \mathbb{T}_k.$$

An important relation between the generalized exponential function and the monomials is given by $\hat{e}_\lambda(t, s) = \sum_{k=0}^{\infty} \lambda^k \hat{h}_k(t, s)$ for $s, t \in \mathbb{T}$ with $t \geq s$, where $\lambda \in \mathcal{R}_c(\mathbb{T})$.

Definition 2.1. The nabla new integral transform ([6]-[8]) of a function $f: \mathbb{T} \rightarrow \mathbb{R}$ is defined by

$$K_\nabla\{f\}(z, v) := \frac{1}{v} \int_v^\infty \hat{e}_{\ominus_v \frac{1}{v^2}}(\rho(\tau), z) f(\tau) \nabla\tau \text{ for } z \in D,$$

where D consists of all complex numbers $z \in \mathbb{C}$ for which the improper ∇ -integral exists.

Definition 2.2. The convolution of two functions $f, g: \mathbb{T} \rightarrow \mathbb{R}$ is defined by

$$(f * g)(t) := \int_a^t \hat{f}(t, \rho(s))g(s)\nabla s, \quad t \in \mathbb{T},$$

where \hat{f} is the shift of f introduced in [3]. Then

$$K_{\nabla}\{f * g\}(z, v) = z \cdot K_{\nabla}\{f\}(z, v) \cdot K_{\nabla}\{g\}(z, v).$$

The following result is the nabla version of the result obtained in [4] (change of integration order).

Let $f \in C_{ld}(\mathbb{T}^2)$, then

$$\int_s^t \int_s^\eta f(\eta, \zeta) \nabla \zeta \nabla \eta = \int_s^t \int_{\rho(\zeta)}^t f(\eta, \zeta) \nabla \eta \nabla \zeta \quad \text{for } s, t \in \mathbb{T}.$$

3 (q, h) – Fractional Calculus

Consider the following (q, h) –time scale:

$$\mathbb{T}_{(q,h)}^{t_0} = \{t_0 q^k + [k]_q h: k \in \mathbb{Z}\} \cup \left\{ \frac{h}{1-q} \right\},$$

for $t_0 > 0, q \geq 1, h \geq 0$ and $q + h > 1$. Note that if $q = 1$, then the cluster point $h/(1-q) = -\infty$ is not involved in \mathbb{T} . The forward and backward jump operator is the linear function $\sigma(t) = qt + h$ and $\rho(t) = q^{-1}(t - h)$, respectively. Similarly, the forward and backward graininess is given by $\mu(t) = (q - 1)t + h$ and $\nu(t) = q^{-1}\mu(t)$, respectively. Observe that

$$\sigma^k(t) = q^k t + [k]_q h \quad \text{and} \quad \rho^k(t) = q^{-k}(t - [k]_q h).$$

The following relation $\nu(\rho^k(t)) = q^{-k}\nu(t)$ holds for $t \in \mathbb{T}$. The nabla (q, h) –derivative of the function $f: \mathbb{T}_{(q,h)}^{t_0} \rightarrow \mathbb{R}$ is defined by

$$\nabla_{(q,h)} f(t) := \frac{f(t) - f(\rho(t))}{\nu(t)} = \frac{f(t) - f(\tilde{q}(t - h))}{(1 - \tilde{q})t + \tilde{q}h},$$

where $\tilde{q} = q^{-1}$. Let $t, a \in \mathbb{T}_{(q,h)}^{t_0}$ such that $h/(1-q) \leq a \leq t$ and $f: \mathbb{T} \rightarrow \mathbb{R}$. Then the nabla (q, h) –integral exists and can be calculated (provided $t > a$) via the formula

$${}_a \nabla_{(q,h)}^{-1} f(t) := \int_a^t f(\tau) \nabla \tau = ((1 - q^{-1})t + q^{-1}h) \sum_{k=0}^{n-1} q^{-k} f(q^{-k}t + [-k]_q h).$$

The Taylor monomials and the power functions on $\mathbb{T}_{(q,h)}^{t_0}$ have the forms

$$\hat{h}_n(t, s) = \frac{\prod_{j=0}^{n-1} (\sigma^j(t) - s)}{[n]_q!} = \frac{\prod_{j=0}^{n-1} (t - \rho^j(s))}{[n]_q!} \quad \text{where } (t - s)_{(q,h)}^{(n)} = \prod_{j=0}^{n-1} (t - \rho^j(s)).$$

Respectively, and the extension of the monomials $\hat{h}_n(t, s)$ corresponding to $\mathbb{T}_{(q,h)}^{t_0}$ takes the form

$$\hat{h}_\alpha(t, s) = \frac{(t - s)_{(q,h)}^{(\alpha)}}{\Gamma_{\tilde{q}}(\alpha + 1)}, \quad \alpha \in \mathbb{R}.$$

Lemma 3.1. [5] Let $m \in \mathbb{Z}^+, \alpha \in \mathbb{R}, s, t \in \mathbb{T}_{(q,h)}^{t_0}$ and $n \in \mathbb{Z}^+, n \geq m$ be such that $t = \sigma^n(s)$.

Then

$$\nabla_{(q,h)}^m \hat{h}_\alpha(t, s) = \begin{cases} \hat{h}_{\alpha-m}(t, s), & \alpha \notin \{0, 1, \dots, m-1\}, \\ 0, & \alpha \in \{0, 1, \dots, m-1\}. \end{cases} \quad (3.1)$$

Let $\alpha \in \mathbb{T}_{(q,h)}^{t_0}, a > h/(1-q)$ be fixed. We consider the following restricted (q, h) –time scale:

$$\tilde{\mathbb{T}}_{(q,h)}^{\sigma^i(a)} = \{t \in \mathbb{T}, t \geq \sigma^i(a)\}, \quad i = 0, 1, 2, \dots, \quad (3.2)$$

where the symbol σ^i stands for the i th iterate of σ (analogously, we use the symbol ρ^i). Now we can continue with the introduction of (q, h) –fractional integral and (q, h) –fractional derivative of a function $f: \mathbb{T}_{(q,h)}^{\sigma^i(a)} \rightarrow \mathbb{R}$. Let $t \in \mathbb{T}_{(q,h)}^{\sigma^i(a)}$.

Definition 3.2. [5] The nabla (q, h) –fractional integral (in the sense of Riemann-Liouville) of order $\alpha \in \mathbb{R}^+$ over the time scale interval $[a, t] \cap \mathbb{T}_{(q,h)}^a$ is defined by

$${}_a\nabla_{(q,h)}^{-\alpha}f(t) = \int_a^t \hat{h}_{\alpha-1}(t, \rho(\tau))f(\tau)\nabla\tau. \quad (3.3)$$

The nabla (q, h) –fractional derivative (in the sense of Caputo) of order $\alpha \in \mathbb{R}^+$ is defined by

$${}_a^c\nabla_{(q,h)}^\alpha f(t) = {}_a\nabla^{-(m-\alpha)}\nabla^m f(t) = \int_a^t \hat{h}_{m-\alpha-1}(t, \rho(\tau))\nabla^m f(\tau)\nabla\tau, \quad (3.4)$$

where $m \in \mathbb{Z}^+$ is such that $m - 1 < \alpha \leq m$.

4. Application

In this section we give solutions for fractional initial value problems on $a \in \mathbb{T}_{(q,h)}^{t_0}$.

Definition 4.1. Let $\alpha, \beta, \gamma \in \mathbb{R}$. The (q, h) –Mittag-Leffler function $E_{\alpha,\beta}^{s,\lambda}(t)$ is defined by

$$E_{\alpha,\beta}^{s,\lambda}(t) = \sum_{k=0}^{\infty} \lambda^k \hat{h}_{\alpha k + \beta - 1}(t, s) \quad \left(= \sum_{k=0}^{\infty} \lambda^k \frac{(t-s)_{(q,h)}^{(\alpha k + \beta - 1)}}{\Gamma_{\tilde{q}}(\alpha k + \beta)} \right), \quad (4.1)$$

for $s, t \in \mathbb{T}_{(q,h)}^{\sigma(a)}$ and $t \geq s$. It is easy to check that the series on the right-hand side converges (absolutely) if $|\lambda|(v(t))^\alpha < 1$. Now we give (q, h) –new integral transforms of fractional nabla integral and Caputo nabla derivative on $\mathbb{T}_{(q,h)}^{t_0}$. For $t, s \in \mathbb{T}_{(q,h)}^{t_0}$ and $\alpha \in \mathbb{R}^+$, we have

$$\hat{K}_{(q,h)}\{\hat{h}_\alpha(t, v)\}(z) = z^{2\alpha+1}.$$

From the definition of (q, h) –Mittag-Leffler function $E_{\alpha,\beta}^{s,\lambda}(t)$, we have

$$\begin{aligned} \hat{K}_{(q,h)}\{E_{\alpha,\beta}^{a,\lambda}(t)\}(z) &= \hat{K}_{(q,h)}\left\{\sum_{k=0}^{\infty} \lambda^k \hat{h}_{\alpha k + \beta - 1}(\cdot, a)\right\}(z) \\ &= \sum_{k=0}^{\infty} \lambda^k \hat{K}_{(q,h)}\{\hat{h}_{\alpha k + \beta - 1}(\cdot, a)\}(z) = \frac{z^{2\beta-1}}{1-\lambda z^{2\alpha}} \text{ provided } |z^{2\alpha}\lambda| < 1. \end{aligned} \quad (4.2)$$

For $\alpha = 1, \beta = 1$ and $t \in \mathbb{T}_{(q,h)}^{\sigma^i(a)}$, we have

$$E_{1,1}^{a,\lambda}(t) = \hat{e}_\lambda(t, a). \quad (4.3)$$

Hence

$$\hat{K}_{(q,h)}\{\hat{e}_\lambda(t, a)\}(z) = \hat{K}_{(q,h)}\{E_{1,1}^{a,\lambda}(t)\}(z) = \frac{z}{1-\lambda z^2}. \quad (4.4)$$

Theorem 4.2. The (q, h) –new integral transform of fractional nabla integral is given by

$$\hat{K}_{(q,h)}\{{}_a\nabla_{(q,h)}^{-\alpha}f\}(z) = z^{2\alpha} \cdot \hat{K}_{(q,h)}\{f\}(z). \quad (4.5)$$

Proof. By convolution, one may write

$${}_a\nabla_{(q,h)}^{-\alpha}f(t) = \int_a^t \hat{h}_{\alpha-1}(t, \rho(\tau))f(\tau)\nabla\tau = (\hat{h}_{\alpha-1}(\cdot, a) * f)(t). \quad (4.6)$$

Thus

$$\widehat{K}_{(q,h)}\{{}_a\nabla_{(q,h)}^{-\alpha}f\}(z) = z \cdot \widehat{K}_{(q,h)}\{\hat{h}_{\alpha-1}(t, a)\}(z) \cdot \widehat{K}_{(q,h)}\{f\}(z) = z^{2\alpha} \cdot \widehat{K}_{(q,h)}\{f\}(z).$$

Theorem 4.3. The (q, h) –new integral transform of Caputo nabla derivative is given by

$$\widehat{K}_{(q,h)}\{{}_a^c\nabla_{(q,h)}^{\alpha}f\}(z) = \frac{\widehat{K}_{(q,h)}\{f\}(z)}{z^{2\alpha}} - \sum_{k=0}^{m-1} \frac{\nabla_{(q,h)}^k f(a)}{z^{2(\alpha-k)-1}}, \quad (4.7)$$

where $m - 1 < \alpha \leq m$. In particular, if $0 < \alpha \leq 1$, then

$$\widehat{K}_{(q,h)}\{{}_a^c\nabla_{(q,h)}^{\alpha}f\}(z) = \frac{\widehat{K}_{(q,h)}\{f\}(z)}{z^{2\alpha}} - \frac{f(a)}{z^{2\alpha-1}}, \quad (4.8)$$

where $\nabla_{(q,h)}^0 f(a) = f(a)$, $k = 0$ and $m = 1$. As an application, we apply (q, h) –new integral transform method to derive explicit solutions to the homogeneous equations of the form

$${}_a^c\nabla_{(q,h)}^{\alpha}y(t) - \lambda y(t) = 0, \quad d_k = \nabla^k y(a), \quad k = 0, \dots, m - 1, \quad (4.9)$$

where $t \in \widetilde{\mathbb{T}}_{(q,h)}^{\sigma m(a)}$ ($m \in \mathbb{N}$), $m - 1 < \alpha \leq m$, $\lambda \in \mathbb{R}$, in terms of the (q, h) –Mittag-Leffler functions. The following statement holds.

Theorem 4.4. Let $m \in \mathbb{N}$ be given by $m - 1 < \alpha \leq m$ and $\lambda \in \mathbb{R}$. Then the functions

$$y_i(t) = E_{\alpha, i+1}^{a, \lambda}(t), \quad i = 0, 1, \dots, m - 1, \quad (4.10)$$

yield the fundamental system of solutions to equation (4.9).

Proof. Applying (q, h) –new integral transform to (4.9) and taking (4.7) into account, we have

$$\widehat{K}_{(q,h)}\{y\}(z) = \sum_{i=0}^{m-1} d_i \frac{z^{2i+1}}{1 - \lambda z^{2\alpha}}. \quad (4.11)$$

Formula (4.2) with $\beta = i + 1$ yields

$$\widehat{K}_{(q,h)}\{E_{\alpha, i+1}^{a, \lambda}(t)\}(z) = \frac{z^{2i+1}}{1 - \lambda z^{2\alpha}}. \quad (4.12)$$

Thus, relation (4.11) yields

$$y(t) = \sum_{i=0}^{m-1} d_i y_i(t), \quad \text{where } y_i(t) = E_{\alpha, i+1}^{a, \lambda}(t). \quad \blacksquare$$

Acknowledgments. Express our sincere gratitude to Associate Professor Akli Fundo, whose suggestion led to conduction of this research.

References

1. M.R.S. Rahmat, M.S.M. Noorani: Caputo type fractional difference operator and its application on discrete time scales, *Advanced in Difference Equations*, 2015:160 DOI 10.1186/s13662-015-0496-5, (2015).
2. B.M. Peterson, A (eds.): *Advanced in Dynamic Equations on Time Scales*, Birkhäuser, Boston, (2003).
3. Kisela, T: Power functions and essentials of fractional calculus on isolated time scales, *Adv. Differ. Equ.*, Article ID 259, (2013).
4. Karpuz: Volterra theory on time scales, *Results Math.*, 65, 263-292, (2014).
5. Cermak, J, Kisela, T, Nechvatal, L: Discrete Mittag-Leffler functions in linear fractional Difference equations, *Abstr. Appl. Anal.*, Article ID 565067, (2011).
6. A. Kashuri, A. Fundo: A New Integral Transform, *Advanced in Theoretical and Applied Mathematics*, ISSN 0973-4554, Vol. 8, No. 1, 27-43, (2013).
7. A. Kashuri, A. Fundo, M. Kreku: Mixture of a New Integral Transform and Homotopy Perturbation Method for Solving Nonlinear Partial Differential Equations, *Advances in Pure Mathematics*, Vol. 3, No. 3, 317-323, (2013).
8. A. Kashuri, A. Fundo, R. Liko: New Integral Transform for Solving Some Fractional Differential Equations, *International Journal of Pure and Applied Mathematics*, ISSN 1311-8080, Vol. 103, No. 4, 675-682, (2015).